

## 300 PRO LA®

### Drug:

Oxytetracycline 300mg/mL

Uses: Pneumonia, Bovine Respiratory Disease, Pink Eye, Foot-rot, Bacterial Enteritis, Leptospirosis, Diphtheria, Wooden Tongue

### Dosage:

**Single dose therapy:** Administer 9-13.6mg/lb. of bodyweight once as a single injection for the control and treatment of Pneumonia and Bovine Respiratory Disease and treatment of Pink Eye.

**Multi-dose therapy:** For other indications administer 5mg/ lb. Bodyweight once every 24 hours.

### Frequency:

**Single dose therapy:** Administer as a one-time injection at a rate of 9-13.6 mg/ lb. animal bodyweight

**Multi-dose therapy:** Administer once every 24 hours at a rate of 5mg/lb. animal bodyweight.

### Route:

This drug is labeled for Intramuscular (IM) or Subcutaneous (SQ) injection. Do not administer intramuscularly in the neck of small calves due to lack of sufficient muscle mass.

### Duration:

Treatment should not to exceed a total of four (4) consecutive days. If improvement is not noted within 24 to 48 hours of the beginning of treatment, diagnosis and therapy should be re-evaluated.

### Withhold:

Milk **NOT FOR USE IN LACTATING DAIRY ANIMALS**

Meat **28 Days**