

# Baytril

## Drug:

Enrofloxacin

Uses: **NOT FOR DAIRY ANIMALS OLDER THAN 20 MONTHS or CALVES INTENDED FOR VEAL**

Bovine Respiratory Disease (BRD) complex associated with Mannheimia haemolytica, Pasteurella multocida and Histophilus somni in beef and non-lactating dairy cattle.

## Dosage:

Baytril® allows for flexible dosing in cattle.

**Single-Dose Therapy:** Administer once, 3.4-5.7 mL/100 lbs. of animal body weight.

**Multiple-Day Therapy:** Administer daily, 1.1-2.3 mL/100 lbs. animal body weight.

## Frequency:

**Single-Dose Therapy:** Administer once

**Multiple Day Therapy:** Administer once daily (every 24 hours).

## Route:

This drug is labeled for Subcutaneous (SQ) injection. No more than 10mL should be injected into any single location. For example, a 20mL dose should be injected into two separate locations, preferably in the neck to insure minimal tissue damage.

## Duration:

Baytril® should be administered for no less than 3 days, but not more than 5 consecutive days.

## Withhold:

Milk **NOT FOR DAIRY ANIMALS OLDER THAN 20 MONTHS, LACTATING DAIRY CATTLE or CALVES INTENDED FOR VEAL.**

Meat **28 Days**

## Special Considerations:

There are no provisions under the Animal Medicinal Drug Use Clarification Act of 1994 (AMDUCA) for the extra-label use of Baytril®. Thus, any administration of Baytril®, for an indication other than those specifically described on the label, constitutes a violation of federal law.