

Micotil

Drug:

300mg tilmicosin phosphate, USP per mL

Uses: **NOT FOR DAIRY ANIMALS OLDER THAN 20 MONTHS or IN ANIMALS INTENDED FOR HUMAN CONSUMPTION WITHIN 42 DAYS OF INJECTION**

Micotil is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* and for the treatment of ovine respiratory disease (ORD) associated with *Mannheimia haemolytica*. Micotil is indicated for the control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia haemolytica*.

Dosage:

In cattle, administer a single dose of 1.5 to 3 mL per 100 lbs.

In sheep greater than 30lbs, administer a single dose of 1.5 mL per 100 lbs.

Frequency:

Once

Route:

This drug is labeled for Subcutaneous (SQ) injection. No more than 10mL should be injected into any single location. For example, a 20mL dose should be injected into two separate locations, preferably in the neck to insure minimal tissue damage.

Duration:

This product is designed as a single dose therapy and should not be repeated without consent of a licensed veterinarian.

Withhold:

Milk **NOT APPROVED FOR LACTATING DAIRY CATTLE**

Meat **42 Days**

Special Considerations:

Human Warning: Injection of this drug in humans has been associated with fatalities. Keep out of reach of children. Do not use in automatically powered syringes. Exercise **extreme caution** to avoid accidental self-injection. In case of human injection, consult a physician immediately and apply ice or cold pack to injection site while avoiding direct contact with the skin. Emergency medical telephone numbers are 1-800-722-0987 or 1-800-428-4441. Avoid contact with eyes and oral cavity.