

Nuflor®

Drug:

Florfenicol 300mg/mL

Uses: **NOT FOR DAIRY ANIMALS OLDER THAN 20 MONTHS, LACTATING DAIRY CATTLE or CALVES INTENDED FOR VEAL**

Bovine Respiratory Disease

Dosage:

Administer 2 doses at a rate of 3mL Nuflor®/100lbs of animal body weight 48 hours apart. Alternatively, a single dose regimen can be followed with a one time injection at a dose rate of 6mL Nuflor®/100lbs of animal body weight.

Frequency:

Nuflor® is labeled as a 2-dose regimen 48 hours apart or a single dose-one time dosing regimen.

Route:

This drug is labeled for Intramuscular (IM) injection at the 3mL/100lb dosage and Subcutaneous (SQ) injection at the 6mL/100lbs dosage. No more than 10mL should be injected into any single location. For example, a 20mL dose should be injected into two separate locations, preferably in the neck to insure minimal tissue damage.

Duration:

This drug should be administered for not less than 4 days and no longer than 7 days, unless directed by a veterinarian.

Withhold:

Milk **NOT FOR DAIRY ANIMALS OLDER THAN 20 MONTHS, LACTATING DAIRY CATTLE or CALVES INTENDED FOR VEAL**

Meat **28 days for IM injection at 3mL/100lbs
38 days for SQ injection at 6mL/100lbs**